

The Strong Onsager Conjecture

Matt Novack
joint with Hyunju Kwon and Vikram Giri

Purdue University

Summer School: Deterministic and Random Features of Fluids
July 4th, 2023

Navier-Stokes and Euler Equations

$$\begin{aligned}\partial_t u + \operatorname{div}(u \otimes u) + \nabla p - \nu \Delta u &= f \\ \operatorname{div} u &= 0\end{aligned}$$

- $u(t, \cdot) : \mathbb{T}^3 \rightarrow \mathbb{R}^3$, $p(t, \cdot) : \mathbb{T}^3 \rightarrow \mathbb{R}$, $f(t, \cdot) : \mathbb{T}^3 \rightarrow \mathbb{R}^3$
- NSE - $\nu > 0$, Euler - $\nu = 0$

Navier-Stokes and Euler Equations

$$\begin{aligned}\partial_t u + \operatorname{div}(u \otimes u) + \nabla p - \nu \Delta u &= f \\ \operatorname{div} u &= 0\end{aligned}$$

- $u(t, \cdot) : \mathbb{T}^3 \rightarrow \mathbb{R}^3$, $p(t, \cdot) : \mathbb{T}^3 \rightarrow \mathbb{R}$, $f(t, \cdot) : \mathbb{T}^3 \rightarrow \mathbb{R}^3$
- NSE - $\nu > 0$, Euler - $\nu = 0$
- **Focus:** Turbulent regime $\nu \rightarrow 0$
- **Facts:** (1) Anomalous dissipation of energy, (2) $4/5$ -law, (3) intermittency

Navier-Stokes and Euler Equations

$$\begin{aligned}\partial_t u + \operatorname{div}(u \otimes u) + \nabla p - \nu \Delta u &= f \\ \operatorname{div} u &= 0\end{aligned}$$

- $u(t, \cdot) : \mathbb{T}^3 \rightarrow \mathbb{R}^3$, $p(t, \cdot) : \mathbb{T}^3 \rightarrow \mathbb{R}$, $f(t, \cdot) : \mathbb{T}^3 \rightarrow \mathbb{R}^3$
- NSE - $\nu > 0$, Euler - $\nu = 0$
- **Focus:** Turbulent regime $\nu \rightarrow 0$
- **Facts:** (1) Anomalous dissipation of energy, (2) $4/5$ -law, (3) intermittency
- **Onsager program:** Build solutions to the PDEs consistent with experiments and numerics!

Main Theorem

Theorem (Giri-Kwon-N., '23)

For any fixed $\beta < 1/3$, there exist weak solutions to the 3D Euler equations

$$\begin{aligned}\partial_t u + \operatorname{div}(u \otimes u) + \nabla p &= 0 \\ \operatorname{div} u &= 0\end{aligned}$$

which, in addition, belong to $C_t^0 B_{3,\infty}^\beta(\mathbb{T}^3)$ and satisfy the local energy inequality

$$\partial_t \left(\frac{1}{2} |u|^2 \right) + \operatorname{div} \left(u \left(\frac{1}{2} |u|^2 + p \right) \right) \leq 0$$

in the sense of distributions.

Turbulence Basics

- Navier-Stokes equations for an incompressible fluid of constant density

$$\begin{aligned}\partial_t u + \operatorname{div}(u \otimes u) &= \frac{1}{\operatorname{Re}} \Delta u - \nabla p + f \\ \operatorname{div} u &= 0\end{aligned}$$

u is velocity, p is pressure, f is an external force


- The *Reynolds number*

$$\operatorname{Re} = \frac{UL}{\nu} = \frac{(\text{characteristic velocity}) \cdot (\text{characteristic length})}{\text{kinematic viscosity}}$$

- Euler equations correspond to $\operatorname{Re} = \infty$, or $\nu = 0$

Turbulence Basics

What happens as the Reynolds number increases?



Flow behind a cylinder at $Re = 1.54$

Turbulence Basics


What happens as the Reynolds number increases?



Flow behind a cylinder at $Re = 140$

Turbulence Basics


What happens as the Reynolds number increases?



Flow behind a grid at $Re = 1800$

Turbulence Basics

- *Homogeneous isotropic turbulence* arises at large Reynolds numbers (or small ν)
- What about anomalous dissipation, the $4/5$ law, and intermittency?



Contour plot of dissipation in a
turbulent velocity field

Source: Kaneda-Ishihara '05

Fact #1: Anomalous Dissipation

$$\partial_t u^\nu + (u^\nu \cdot \nabla) u^\nu = \nu \Delta u^\nu - \nabla p^\nu, \quad \operatorname{div} u^\nu = 0$$

- Pointwise energy balance for smooth solutions

$$\partial_t \left(\frac{1}{2} |u^\nu|^2 \right) + \operatorname{div} \left(\left(\frac{1}{2} |u^\nu|^2 + p^\nu \right) u^\nu - \nu \nabla \frac{|u^\nu|^2}{2} \right) = -\nu |\nabla u^\nu|^2$$

Fact #1: Anomalous Dissipation

$$\partial_t u^\nu + (u^\nu \cdot \nabla) u^\nu = \nu \Delta u^\nu - \nabla p^\nu, \quad \operatorname{div} u^\nu = 0$$

- Pointwise energy balance for smooth solutions

$$\partial_t \left(\frac{1}{2} |u^\nu|^2 \right) + \operatorname{div} \left(\left(\frac{1}{2} |u^\nu|^2 + p^\nu \right) u^\nu - \nu \nabla \frac{|u^\nu|^2}{2} \right) = -\nu |\nabla u^\nu|^2$$

- Integrating in \mathbb{T}^3 and from 0 to T , we have

$$\frac{1}{2} \|u^\nu(T, \cdot)\|_{L^2(\mathbb{T}^3)}^2 - \frac{1}{2} \|u^\nu(0, \cdot)\|_{L^2(\mathbb{T}^3)}^2 = - \int_0^T \nu \|\nabla u^\nu(t, \cdot)\|_{L^2(\mathbb{T}^3)}^2 dt$$

Fact #1: Anomalous Dissipation

$$\partial_t u^\nu + (u^\nu \cdot \nabla) u^\nu = \nu \Delta u^\nu - \nabla p^\nu, \quad \operatorname{div} u^\nu = 0$$

- Pointwise energy balance for smooth solutions

$$\partial_t \left(\frac{1}{2} |u^\nu|^2 \right) + \operatorname{div} \left(\left(\frac{1}{2} |u^\nu|^2 + p^\nu \right) u^\nu - \nu \nabla \frac{|u^\nu|^2}{2} \right) = -\nu |\nabla u^\nu|^2$$

- Integrating in \mathbb{T}^3 and from 0 to T , we have

$$\frac{1}{2} \|u^\nu(T, \cdot)\|_{L^2(\mathbb{T}^3)}^2 - \frac{1}{2} \|u^\nu(0, \cdot)\|_{L^2(\mathbb{T}^3)}^2 = - \int_0^T \nu \|\nabla u^\nu(t, \cdot)\|_{L^2(\mathbb{T}^3)}^2 dt$$

- Thus smooth Euler solutions conserve energy, and dissipation in smooth Navier-Stokes solutions is caused by $\nu \Delta u^\nu$

Fact #1: Anomalous Dissipation

$$\partial_t u^\nu + (u^\nu \cdot \nabla) u^\nu = \nu \Delta u^\nu - \nabla p^\nu, \quad \operatorname{div} u^\nu = 0$$

- Energy balance for weak solutions (obtained by mollifying and passing to the limit)

$$\partial_t \left(\frac{1}{2} |u^\nu|^2 \right) + \operatorname{div} \left(\left(\frac{1}{2} |u^\nu|^2 + p^\nu \right) u^\nu - \nu \nabla \frac{|u^\nu|^2}{2} \right) + \nu |\nabla u^\nu|^2 + D[u^\nu] = 0$$

Fact #1: Anomalous Dissipation

$$\partial_t u^\nu + (u^\nu \cdot \nabla) u^\nu = \nu \Delta u^\nu - \nabla p^\nu, \quad \operatorname{div} u^\nu = 0$$

- Energy balance for weak solutions (obtained by mollifying and passing to the limit)

$$\partial_t \left(\frac{1}{2} |u^\nu|^2 \right) + \operatorname{div} \left(\left(\frac{1}{2} |u^\nu|^2 + p^\nu \right) u^\nu - \nu \nabla \frac{|u^\nu|^2}{2} \right) + \nu |\nabla u^\nu|^2 + D[u^\nu] = 0$$

- The nonlinearity contributes the *Duchon-Robert measure*

$$D[u^\nu](t, x) = \lim_{\ell \rightarrow 0} \frac{1}{4} \int_{\mathbb{T}^3} \nabla \phi_\ell(z) \cdot (u(t, x+z) - u(t, x)) |u(t, x+z) - u(t, x)|^2 dz$$

Fact #1: Anomalous Dissipation

$$\partial_t u^\nu + (u^\nu \cdot \nabla) u^\nu = \nu \Delta u^\nu - \nabla p^\nu, \quad \operatorname{div} u^\nu = 0$$

- Energy balance for weak solutions (obtained by mollifying and passing to the limit)

$$\partial_t \left(\frac{1}{2} |u^\nu|^2 \right) + \operatorname{div} \left(\left(\frac{1}{2} |u^\nu|^2 + p^\nu \right) u^\nu - \nu \nabla \frac{|u^\nu|^2}{2} \right) + \nu |\nabla u^\nu|^2 + D[u^\nu] = 0$$

- The nonlinearity contributes the *Duchon-Robert measure*

$$D[u^\nu](t, x) = \lim_{\ell \rightarrow 0} \frac{1}{4} \int_{\mathbb{T}^3} \nabla \phi_\ell(z) \cdot (u(t, x+z) - u(t, x)) |u(t, x+z) - u(t, x)|^2 dz$$

- Zeroth law of turbulence (no proof exists!)

$$\varepsilon = \liminf_{\nu \rightarrow 0} \underbrace{\left\langle \nu |\nabla u^\nu|^2 + D[u^\nu] \right\rangle}_{\varepsilon^\nu} > 0$$

Fact #1: Anomalous Dissipation

- Caffarelli-Kohn-Nirenberg's "suitable solutions" to Navier-Stokes satisfy

$$u^\nu \in L_t^\infty L_x^2 \cap L_t^2 W_x^{1,2}, \quad D[u^\nu] \geq 0$$

- If suitable solutions u^ν converge in $L_{t,x}^3$ to an Euler solution u , then $D[u] \geq 0$

Fact #1: Anomalous Dissipation

- Caffarelli-Kohn-Nirenberg's "suitable solutions" to Navier-Stokes satisfy

$$u^\nu \in L_t^\infty L_x^2 \cap L_t^2 W_x^{1,2}, \quad D[u^\nu] \geq 0$$

- If suitable solutions u^ν converge in $L_{t,x}^3$ to an Euler solution u , then $D[u] \geq 0$
- Notice that $D[u^\nu]$, given by

$$D[u^\nu] = \lim_{\ell \rightarrow 0} \frac{1}{4} \int_{\mathbb{T}^3} \nabla \phi_\ell(z) \cdot (u^\nu(t, x+z) - u^\nu(t, x)) |u^\nu(t, x+z) - u^\nu(t, x)|^2 dz,$$

has units U^3/L

Fact #1: Anomalous Dissipation

- Caffarelli-Kohn-Nirenberg's "suitable solutions" to Navier-Stokes satisfy

$$u^\nu \in L_t^\infty L_x^2 \cap L_t^2 W_x^{1,2}, \quad D[u^\nu] \geq 0$$

- If suitable solutions u^ν converge in $L_{t,x}^3$ to an Euler solution u , then $D[u] \geq 0$
- Notice that $D[u^\nu]$, given by

$$D[u^\nu] = \lim_{\ell \rightarrow 0} \frac{1}{4} \int_{\mathbb{T}^3} \nabla \phi_\ell(z) \cdot (u^\nu(t, x+z) - u^\nu(t, x)) |u^\nu(t, x+z) - u^\nu(t, x)|^2 dz,$$

has units U^3/L

- Thus if $D[u] \neq 0$, u^ν cannot remain bounded in $L_t^3 B_{3,\infty,x}^\alpha$ for $\alpha > 1/3$ as $\nu \rightarrow 0$, where

$$\|f\|_{B_{3,\infty}^\alpha(\mathbb{T}^3)} = \sup_{|z|>0} \frac{1}{|z|^\alpha} \|f(\cdot+z) - f(\cdot)\|_{L^3(\mathbb{T}^3)}$$

Fact #2: Kolmogorov's $4/5$ law

- **K41 Assumptions:** the zeroth law ($\varepsilon > 0$), translation, rotation, and scaling symmetries for law of $u^\nu(t, x + \ell \hat{z}) - u^\nu(t, x)$ (here $\ell > 0, \hat{z} \in \mathbb{S}^2$)

Fact #2: Kolmogorov's $4/5$ law

- **K41 Assumptions:** the zeroth law ($\varepsilon > 0$), translation, rotation, and scaling symmetries for law of $u^\nu(t, x + \ell \hat{z}) - u^\nu(t, x)$ (here $\ell > 0, \hat{z} \in \mathbb{S}^2$)
- **K41 Claims:** longitudinal structure functions satisfy

$$S_p^\parallel(\ell) = \langle ((u^\nu(t, x + \ell \hat{z}) - u^\nu(t, x)) \cdot \hat{z})^p \rangle \approx (\varepsilon \ell)^{p/3}$$

Fact #2: Kolmogorov's $4/5$ law

- **K41 Assumptions:** the zeroth law ($\varepsilon > 0$), translation, rotation, and scaling symmetries for law of $u^\nu(t, x + \ell \hat{z}) - u^\nu(t, x)$ (here $\ell > 0, \hat{z} \in \mathbb{S}^2$)

- **K41 Claims:** longitudinal structure functions satisfy

$$S_p^\parallel(\ell) = \langle ((u^\nu(t, x + \ell \hat{z}) - u^\nu(t, x)) \cdot \hat{z})^p \rangle \approx (\varepsilon \ell)^{p/3}$$

- **Regularity:** K41-style scaling suggests that for $p \in [1, \infty)$,

$$\underbrace{\sup_{0 < z \leq 1} |z|^{-\frac{p}{3}} \|u(t, \cdot + z) - u(t, \cdot)\|_{L^p(\mathbb{T}^3)}^p \approx \varepsilon^{p/3}}_{\implies u(t, \cdot) \in B_{p, \infty}^{1/3}} \implies u(t, \cdot) \in C^{1/3}$$

Fact #2: Kolmogorov's $4/5$ law

- **K41 Assumptions:** the zeroth law ($\varepsilon > 0$), translation, rotation, and scaling symmetries for law of $u^\nu(t, x + \ell \hat{z}) - u^\nu(t, x)$ (here $\ell > 0, \hat{z} \in \mathbb{S}^2$)

- **K41 Claims:** longitudinal structure functions satisfy

$$S_p^\parallel(\ell) = \langle ((u^\nu(t, x + \ell \hat{z}) - u^\nu(t, x)) \cdot \hat{z})^p \rangle \approx (\varepsilon \ell)^{p/3}$$

- **Regularity:** K41-style scaling suggests that for $p \in [1, \infty)$,

$$\underbrace{\sup_{0 < z \leq 1} |z|^{-\frac{p}{3}} \|u(t, \cdot + z) - u(t, \cdot)\|_{L^p(\mathbb{T}^3)}^p \approx \varepsilon^{p/3}}_{\implies u(t, \cdot) \in B_{p, \infty}^{1/3}} \implies u(t, \cdot) \in C^{1/3}$$

- Local, deterministic $4/5$ law (Eyink, '02)

$$\begin{aligned} \lim_{\ell \rightarrow 0} \frac{1}{\ell} S_3^\parallel(\ell) &= \lim_{\ell \rightarrow 0} \frac{1}{\ell} \int_{\mathbb{T}^3} \int_{\mathbb{S}^2} [(u^\nu(t, x + \ell z) - u^\nu(t, x)) \cdot z]^3 dz dx \\ &= -\frac{4}{5} D[u^\nu] \end{aligned}$$

Onsager's Conjecture and the $L_t^\infty C_x^{1/3}$ Threshold

- "It is of some interest to note that in principle, turbulent dissipation as described could take place just as readily without the final assistance by viscosity. In the absence of viscosity, the standard proof of the conservation of energy does not apply, because the velocity field does not remain differentiable!" – Onsager, '49

Onsager's Conjecture and the $L_t^\infty C_x^{1/3}$ Threshold

- “It is of some interest to note that in principle, turbulent dissipation as described could take place just as readily without the final assistance by viscosity. In the absence of viscosity, the standard proof of the conservation of energy does not apply, because the velocity field does not remain differentiable!” – Onsager, '49
- Recall that

$$\begin{aligned} & \partial_t \left(\frac{1}{2} |u|^2 \right) (t, x) + \operatorname{div} \left(\left(\frac{1}{2} |u|^2 + p \right) u \right) (t, x) \\ &= - \lim_{\ell \rightarrow 0} \frac{1}{4} \int_{\mathbb{T}^3} \nabla \phi_\ell(z) \cdot (u(t, x+z) - u(t, x)) |u(t, x+z) - u(t, x)|^2 dz \end{aligned}$$

Onsager's Conjecture and the $L_t^\infty C_x^{1/3}$ Threshold

- “It is of some interest to note that in principle, turbulent dissipation as described could take place just as readily without the final assistance by viscosity. In the absence of viscosity, the standard proof of the conservation of energy does not apply, because the velocity field does not remain differentiable!” – Onsager, '49
- Recall that

$$\begin{aligned} & \partial_t \left(\frac{1}{2} |u|^2 \right) (t, x) + \operatorname{div} \left(\left(\frac{1}{2} |u|^2 + p \right) u \right) (t, x) \\ &= - \lim_{\ell \rightarrow 0} \frac{1}{4} \int_{\mathbb{T}^3} \nabla \phi_\ell(z) \cdot (u(t, x+z) - u(t, x)) |u(t, x+z) - u(t, x)|^2 dz \end{aligned}$$

and so conservation of energy follows from $D[u] = 0$, which holds if $u \in L_t^3 B_{3,\infty}^\alpha$ for $\alpha > 1/3$ (Eyink '92, Constantin-E-Titi '94, Duchon-Robert '00)

Onsager's Conjecture and the $L_t^\infty C_x^{1/3}$ Threshold

- “It is of some interest to note that in principle, turbulent dissipation as described could take place just as readily without the final assistance by viscosity. In the absence of viscosity, the standard proof of the conservation of energy does not apply, because the velocity field does not remain differentiable!” – Onsager, '49
- Recall that

$$\begin{aligned} & \partial_t \left(\frac{1}{2} |u|^2 \right) (t, x) + \operatorname{div} \left(\left(\frac{1}{2} |u|^2 + p \right) u \right) (t, x) \\ &= - \lim_{\ell \rightarrow 0} \frac{1}{4} \int_{\mathbb{T}^3} \nabla \phi_\ell(z) \cdot (u(t, x+z) - u(t, x)) |u(t, x+z) - u(t, x)|^2 dz \end{aligned}$$

and so conservation of energy follows from $D[u] = 0$, which holds if $u \in L_t^3 B_{3,\infty}^\alpha$ for $\alpha > 1/3$ (Eyink '92, Constantin-E-Titi '94, Duchon-Robert '00)

- If $\alpha < 1/3$, the kinetic energy of 3D Euler solutions need not be conserved (Isett '18) and can dissipate (Buckmaster-De Lellis-Székelyhidi-Vicol '19)

Adding to the story: local energy inequality and intermittency

- The solutions of Isett and Buckmaster et. al. do not satisfy $D[u] \geq 0$, and so cannot arise as limits of suitable Navier-Stokes solutions

Adding to the story: local energy inequality and intermittency

- The solutions of Isett and Buckmaster et. al. do not satisfy $D[u] \geq 0$, and so cannot arise as limits of suitable Navier-Stokes solutions
- Solutions satisfying $D[u] \geq 0$, also known as the *local energy inequality*

$$\partial_t \left(\frac{1}{2} |u|^2 \right) + \operatorname{div} \left(\left(\frac{1}{2} |u|^2 + p \right) u \right) = -D[u^\nu] \leq 0,$$

have only been shown to exist in $C^{1/7-}$ (De Lellis-Kwon '22, following Isett '22); there are fundamental obstructions to reaching $C^{1/3-}$

Adding to the story: local energy inequality and intermittency


- The solutions of Isett and Buckmaster et. al. do not satisfy $D[u] \geq 0$, and so cannot arise as limits of suitable Navier-Stokes solutions
- Solutions satisfying $D[u] \geq 0$, also known as the *local energy inequality*

$$\partial_t \left(\frac{1}{2} |u|^2 \right) + \operatorname{div} \left(\left(\frac{1}{2} |u|^2 + p \right) u \right) = -D[u^\nu] \leq 0,$$

have only been shown to exist in $C^{1/7-}$ (De Lellis-Kwon '22, following Isett '22); there are fundamental obstructions to reaching $C^{1/3-}$

- Conservation of energy requires only $L_t^3 B_{3,\infty}^\alpha$ for $\alpha > 1/3$, but dissipative solutions belong to $C_{t,x}^\alpha$ for $\alpha < 1/3$... is this merely a curiosity concerning function spaces?

Intermittency: Deviations from K41/Onsager



— = K41

Intermittency: Deviations from K41/Onsager

- **Onsager, unpublished work** - “[Anomalous scaling for ζ_2] would require a “spotty” distribution of the regions in which the velocity varies rapidly”
- **Kolmogorov '62** - “I have formulated appropriate modifications to the two similarity hypotheses that I put forward in 1941 ...”
- **Chen, Dhruva, Kurien, Sreenivasan, Taylor '05** - “It is now believed that the scaling exponents of moments of velocity increments are anomalous ... anomalous scaling is a genuine result worth of a serious theoretical effort.”
- **Iyer, Sreenivasan, Yeung '20** - “The $4/5$ -ths law holds in an intermediate range of scales and the second-order exponent over the same range of scales is *anomalous*, departing from the self-similar value of $2/3$.”
- **See also** - Ishihara-Kaneda-Gotoh, Frisch, Anselmet-Gagne-Hopfinger-Antonia, ...


Takeaway: $B_{3,\infty}^{1/3} \cap L^\infty$ may be the correct space

Intermittency: Deviations from K41/Onsager

- **Symmetry assumptions:** Turbulence is *isotropic, homogeneous, but not purely self-similar ... fewer eddies of higher intensity!*

Intermittency: Deviations from K41/Onsager

- **Symmetry assumptions:** Turbulence is *isotropic, homogeneous, but not purely self-similar ... fewer eddies of higher intensity!*
- **Dissipativity assumption:** Dissipation occurs even in the absence of viscosity
- **Implications for regularity:** Cantor function, Heaviside function ($B_{p,\infty}^{1/p}$)



Strong Onsager Conjectures

Consider weak solutions u to the Euler equations, with the local energy balance

$$\begin{cases} \partial_t u + \operatorname{div}(u \otimes u) + \nabla p = 0 \\ \operatorname{div} u = 0 \\ \partial_t \left(\frac{1}{2}|u|^2\right) + \operatorname{div} \left(\left(\frac{1}{2}|u|^2 + p\right)u\right) + D[u] = 0. \end{cases}$$

Strong Onsager Conjectures

Consider weak solutions u to the Euler equations, with the local energy balance

$$\begin{cases} \partial_t u + \operatorname{div}(u \otimes u) + \nabla p = 0 \\ \operatorname{div} u = 0 \\ \partial_t \left(\frac{1}{2}|u|^2\right) + \operatorname{div} \left(\left(\frac{1}{2}|u|^2 + p\right)u\right) + D[u] = 0. \end{cases}$$

Conjecture (Strong L^3 Onsager Conjecture)

For any $\alpha < 1/3$, there exist weak solutions $u \in L_t^\infty B_{3,\infty}^\alpha$ of 3D Euler for which $D[u] \geq 0$ and $\frac{d}{dt} \|u\|_{L^2}^2 < 0$.

The Strong L^3 Onsager Theorem (and its antecedents)

Theorem (Buckmaster-Masmoudi-N.-Vicol, '21)

There exist solutions to 3D Euler which have decreasing kinetic energy and belong to $L_t^\infty H_x^{1/2-}$ (intermittent, but no $4/5$ law or local energy inequality)

The Strong L^3 Onsager Theorem (and its antecedents)

Theorem (Buckmaster-Masmoudi-N.-Vicol, '21)

There exist solutions to 3D Euler which have decreasing kinetic energy and belong to $L_t^\infty H_x^{1/2-}$ (intermittent, but no $4/5$ law or local energy inequality)

Theorem (N.-Vicol, '22)

There exist solutions to 3D Euler which have decreasing kinetic energy and belong to $L_t^\infty \left(H_x^{1/2-} \cap L_x^\infty \cap B_{3,\infty}^{1/3-} \right)$ (intermittent, $4/5$ law, but no local energy inequality)

The Strong L^3 Onsager Theorem (and its antecedents)

Theorem (Buckmaster-Masmoudi-N.-Vicol, '21)

There exist solutions to 3D Euler which have decreasing kinetic energy and belong to $L_t^\infty H_x^{1/2-}$ (intermittent, but no $4/5$ law or local energy inequality)

Theorem (N.-Vicol, '22)

There exist solutions to 3D Euler which have decreasing kinetic energy and belong to $L_t^\infty \left(H_x^{1/2-} \cap L_x^\infty \cap B_{3,\infty}^{1/3-} \right)$ (intermittent, $4/5$ law, but no local energy inequality)

- Giri-Kwon-N., '23a - Reprove the results in NV '22 using a “wavelet-inspired” scheme, along with further improvements

The Strong L^3 Onsager Theorem (and its antecedents)

Theorem (Buckmaster-Masmoudi-N.-Vicol, '21)

There exist solutions to 3D Euler which have decreasing kinetic energy and belong to $L_t^\infty H_x^{1/2-}$ (intermittent, but no $4/5$ law or local energy inequality)

Theorem (N.-Vicol, '22)

There exist solutions to 3D Euler which have decreasing kinetic energy and belong to $L_t^\infty \left(H_x^{1/2-} \cap L_x^\infty \cap B_{3,\infty}^{1/3-} \right)$ (intermittent, $4/5$ law, but no local energy inequality)

- Giri-Kwon-N., '23a - Reprove the results in NV '22 using a “wavelet-inspired” scheme, along with further improvements

Theorem (Giri-Kwon-N., '23b)

There exist solutions to 3D Euler which have decreasing kinetic energy, belong to $L_t^\infty \left(L_x^\infty \cap B_{3,\infty}^{1/3-} \right)$, and satisfy $D[u] \geq 0$ (intermittent, $4/5$ law, local energy inequality)

A bird's-eye view of our construction

Basic Strategy: Construct a weak solution $u = \lim_{q \rightarrow \infty} u_q$ in $L_t^\infty \left(L^{\infty-} \cap B_{3,\infty}^{1/3-} \right)$ via Nash iteration

A bird's-eye view of our construction

Basic Strategy: Construct a weak solution $u = \lim_{q \rightarrow \infty} u_q$ in $L_t^\infty \left(L^{\infty-} \cap B_{3,\infty}^{1/3-} \right)$ via Nash iteration

- **Wavelet-inspired scheme:** Replace partial Fourier sums u_q of solution u (i.e. pure frequency decompositions), with partial wavelet decompositions u_q (mixed space-time and frequency decompositions)

A bird's-eye view of our construction

Basic Strategy: Construct a weak solution $u = \lim_{q \rightarrow \infty} u_q$ in $L_t^\infty \left(L^{\infty-} \cap B_{3,\infty}^{1/3-} \right)$ via Nash iteration

- **Wavelet-inspired scheme:** Replace partial Fourier sums u_q of solution u (i.e. pure frequency decompositions), with partial wavelet decompositions u_q (mixed space-time and frequency decompositions)
- **L^3 iteration:** Replace Nash iterations formulated in L^2 or L^∞ -based Sobolev spaces with an L^3 -based framework

A bird's-eye view of our construction

Basic Strategy: Construct a weak solution $u = \lim_{q \rightarrow \infty} u_q$ in $L_t^\infty \left(L^{\infty-} \cap B_{3,\infty}^{1/3-} \right)$ via Nash iteration

- **Wavelet-inspired scheme:** Replace partial Fourier sums u_q of solution u (i.e. pure frequency decompositions), with partial wavelet decompositions u_q (mixed space-time and frequency decompositions)
- **L^3 iteration:** Replace Nash iterations formulated in L^2 or L^∞ -based Sobolev spaces with an L^3 -based framework
- **Intermittent pressure π_q :** Control u_q (and any other important functions) in terms of π_q , and propagate scaling laws comparing different terms in the wavelet expansion of π_q

A bird's-eye view of our construction

Basic Strategy: Construct a weak solution $u = \lim_{q \rightarrow \infty} u_q$ in $L_t^\infty \left(L^{\infty-} \cap B_{3,\infty}^{1/3-} \right)$ via Nash iteration

- **Wavelet-inspired scheme:** Replace partial Fourier sums u_q of solution u (i.e. pure frequency decompositions), with partial wavelet decompositions u_q (mixed space-time and frequency decompositions)
- **L^3 iteration:** Replace Nash iterations formulated in L^2 or L^∞ -based Sobolev spaces with an L^3 -based framework
- **Intermittent pressure π_q :** Control u_q (and any other important functions) in terms of π_q , and propagate scaling laws comparing different terms in the wavelet expansion of π_q

Two questions: How to set up the induction, and how to propagate the inductive assumptions?

Wavelet-inspired, $B_{3,\infty}^{1/3-} \cap L^{\infty-}$ inductive set-up

- Assume the existence of $(u_q, p_q, R_q, \Phi_q, \pi_q)$ satisfying

$$\partial_t u_q + \operatorname{div} (u_q \otimes u_q) + \nabla p_q = \underbrace{\operatorname{div}_x (R_q - \pi_q \operatorname{Id})}_{\rightarrow 0 \text{ as } q \rightarrow \infty}, \quad \operatorname{div} u_q = 0$$

$$\partial_t \left(\frac{|u_q|^2}{2} \right) + \operatorname{div} \left(\left(\frac{|u_q|^2}{2} + p_q \right) u_q \right) \leq \underbrace{\operatorname{div}_{t,x} \Phi_q(\pi_q)}_{\rightarrow 0 \text{ as } q \rightarrow \infty}$$

Wavelet-inspired, $B_{3,\infty}^{1/3-} \cap L^{\infty-}$ inductive set-up

- Assume the existence of $(u_q, p_q, R_q, \Phi_q, \pi_q)$ satisfying

$$\partial_t u_q + \operatorname{div} (u_q \otimes u_q) + \nabla p_q = \underbrace{\operatorname{div}_x (R_q - \pi_q \operatorname{Id})}_{\rightarrow 0 \text{ as } q \rightarrow \infty}, \quad \operatorname{div} u_q = 0$$

$$\partial_t \left(\frac{|u_q|^2}{2} \right) + \operatorname{div} \left(\left(\frac{|u_q|^2}{2} + p_q \right) u_q \right) \leq \underbrace{\operatorname{div}_{t,x} \Phi_q(\pi_q)}_{\rightarrow 0 \text{ as } q \rightarrow \infty}$$

- Heuristics:**

- Let $\lambda_q = a^{(b^q)}$ quantify the inverse of the diameter of an oscillation, and $\lambda_q r_q$ for $r_q \ll 1$ quantify the inverse of the distance between oscillations

Wavelet-inspired, $B_{3,\infty}^{1/3-} \cap L^{\infty-}$ inductive set-up

- Assume the existence of $(u_q, p_q, R_q, \Phi_q, \pi_q)$ satisfying

$$\partial_t u_q + \operatorname{div} (u_q \otimes u_q) + \nabla p_q = \underbrace{\operatorname{div}_x (R_q - \pi_q \operatorname{Id})}_{\rightarrow 0 \text{ as } q \rightarrow \infty}, \quad \operatorname{div} u_q = 0$$

$$\partial_t \left(\frac{|u_q|^2}{2} \right) + \operatorname{div} \left(\left(\frac{|u_q|^2}{2} + p_q \right) u_q \right) \leq \underbrace{\operatorname{div}_{t,x} \Phi_q(\pi_q)}_{\rightarrow 0 \text{ as } q \rightarrow \infty}$$

- Heuristics:**

- Let $\lambda_q = a^{(b^q)}$ quantify the inverse of the diameter of an oscillation, and $\lambda_q r_q$ for $r_q \ll 1$ quantify the inverse of the distance between oscillations

- $u_q = \sum_{q' \leq q} w_{q'}$, where $w_{q'}$ has oscillations of size $\lambda_{q'}^{-1}$, distance $(\lambda_{q'} r_{q'})^{-1}$ between oscillations, and frequency support $[\lambda_{q'} r_{q'}, \lambda_{q'}]$ (not necessarily disjoint from $[\lambda_{q'} r_{q'}, \lambda_{q'}]$!)

Wavelet-inspired, $B_{3,\infty}^{1/3-} \cap L^{\infty-}$ inductive set-up

- Assume the existence of $(u_q, p_q, R_q, \Phi_q, \pi_q)$ satisfying

$$\partial_t u_q + \operatorname{div} (u_q \otimes u_q) + \nabla p_q = \underbrace{\sum_{k=q-\bar{n}+1}^q \operatorname{div}_x (R_q^k - \pi_q^k \operatorname{Id})}_{\rightarrow 0 \text{ as } q \rightarrow \infty}, \quad \operatorname{div} u_q = 0$$

$$\partial_t \left(\frac{|u_q|^2}{2} \right) + \operatorname{div} \left(\left(\frac{|u_q|^2}{2} + p_q \right) u_q \right) \leq \underbrace{\sum_{k=q-\bar{n}+1}^q \operatorname{div}_{t,x} \Phi_q^k(\pi_q^k)}_{\rightarrow 0 \text{ as } q \rightarrow \infty}$$

- Heuristics:**

- Let $\lambda_q \approx a^{(b^q)}$ quantify the inverse of the width of an oscillation, and $\lambda_q r_q$ for $r_q \ll 1$ describe the inverse of the distance between oscillations
- $u_q = \sum_{q' \leq q} w_{q'}$, where $w_{q'}$ has oscillations of size $\lambda_{q'}^{-1}$, distance $(\lambda_{q'} r_{q'})^{-1}$ between oscillations, and frequency support $[\lambda_{q'} r_{q'}, \lambda_{q'}]$
- “Wavelet parameter” \bar{n} quantifies the exchange between space-time and frequency support and decomposes $R_q, \Phi_q,$ and π_q by frequency:

$$\operatorname{supp}_{t,x} w_{q'} \cap w_{q''} = \emptyset \quad \text{if} \quad |q' - q''| < \bar{n}$$

$$\operatorname{supp}_{\xi} \hat{w}_{q'} \cap \hat{w}_{q''} = \emptyset \quad \text{if} \quad |q' - q''| \geq \bar{n}$$

Wavelet-inspired, $B_{3,\infty}^{1/3-} \cap L^{\infty-}$ inductive set-up

- Assume the existence of $(u_q, p_q, R_q, \Phi_q, \pi_q)$ satisfying

$$\partial_t u_q + \operatorname{div} (u_q \otimes u_q) + \nabla p_q = \underbrace{\sum_{k=q-\bar{n}+1}^q \operatorname{div}_x (R_q^k - \pi_q^k \operatorname{Id})}_{\rightarrow 0 \text{ as } q \rightarrow \infty}, \quad \operatorname{div} u_q = 0$$

$$\partial_t \left(\frac{|u_q|^2}{2} \right) + \operatorname{div} \left(\left(\frac{|u_q|^2}{2} + p_q \right) u_q \right) \leq \underbrace{\sum_{k=q-\bar{n}+1}^q \operatorname{div}_{t,x} \Phi_q^k(\pi_q^k)}_{\rightarrow 0 \text{ as } q \rightarrow \infty}$$

- Heuristics:**

- Use parameters λ_q , r_q , and \bar{n} to describe u_q as a “partial wavelet sum,” decompose R_q , Φ_q , π_q by frequency

- Inductive bounds:**

- Assume bounds for $\|\pi\|_{3/2}$, $\|\pi\|_{\infty}$ (which depend on their frequency)

Wavelet-inspired, $B_{3,\infty}^{1/3-} \cap L^{\infty-}$ inductive set-up

- Assume the existence of $(u_q, p_q, R_q, \Phi_q, \pi_q)$ satisfying

$$\partial_t u_q + \operatorname{div} (u_q \otimes u_q) + \nabla p_q = \underbrace{\sum_{k=q-\bar{n}+1}^q \operatorname{div}_x (R_q^k - \pi_q^k \operatorname{Id})}_{\rightarrow 0 \text{ as } q \rightarrow \infty}, \quad \operatorname{div} u_q = 0$$

$$\partial_t \left(\frac{|u_q|^2}{2} \right) + \operatorname{div} \left(\left(\frac{|u_q|^2}{2} + p_q \right) u_q \right) \leq \underbrace{\sum_{k=q-\bar{n}+1}^q \operatorname{div}_{t,x} \Phi_q^k(\pi_q^k)}_{\rightarrow 0 \text{ as } q \rightarrow \infty}$$

- Heuristics:**

- Use parameters λ_q , r_q , and \bar{n} to describe u_q as a “partial wavelet sum,” decompose R_q , Φ_q , π_q by frequency

- Inductive bounds:**

- Assume bounds for $\|\pi\|_{3/2}$, $\|\pi\|_{\infty}$ (which depend on their frequency)
- All other bounds are “pointwise in terms of π ,” e.g.

$$\left| D^N (\partial_t + u \cdot \nabla)^M \pi \right| + \left| D^N (\partial_t + u \cdot \nabla)^M R \right| \lesssim \pi \lambda^N \left(\pi^{1/2} \lambda \right)^M$$

Wavelet-inspired, $B_{3,\infty}^{1/3-} \cap L^{\infty-}$ inductive set-up

- Assume the existence of $(u_q, p_q, R_q, \Phi_q, \pi_q)$ satisfying

$$\partial_t u_q + \operatorname{div} (u_q \otimes u_q) + \nabla p_q = \sum_{k=q-\bar{n}+1}^q \operatorname{div}_x \left(R_q^k - \pi_q^k \operatorname{Id} \right), \quad \operatorname{div} u_q = 0$$

$$\partial_t \left(\frac{|u_q|^2}{2} \right) + \operatorname{div} \left(\left(\frac{|u_q|^2}{2} + p_q \right) u_q \right) \leq \sum_{k=q-\bar{n}+1}^q \operatorname{div}_{t,x} \Phi_q^k(\pi_q^k)$$

- **Heuristics:**

- Use parameters λ_q , r_q , and \bar{n} to decompose u_q into a “partial wavelet sum,”
decompose R_q , Φ_q , π_q by frequency

- **Inductive bounds:**

- Assume bounds for $\|\pi\|_{3/2}$, $\|\pi\|_{\infty}$ (which depend on their frequency)
- All other bounds are *pointwise bounds* in terms of π

- **Scaling law:** Mollifying high-freq. π 's gives rescaled low-freq. π 's

Wavelet-inspired, $B_{3,\infty}^{1/3-} \cap L^{\infty-}$ inductive set-up

- Assume the existence of $(u_q, p_q, R_q, \Phi_q, \pi_q)$ satisfying

$$\partial_t u_q + \operatorname{div} (u_q \otimes u_q) + \nabla p_q = \sum_{k=q-\bar{n}+1}^q \operatorname{div}_x \left(R_q^k - \pi_q^k \operatorname{Id} \right), \quad \operatorname{div} u_q = 0$$

$$\partial_t \left(\frac{|u_q|^2}{2} \right) + \operatorname{div} \left(\left(\frac{|u_q|^2}{2} + p_q \right) u_q \right) \leq \sum_{k=q-\bar{n}+1}^q \operatorname{div}_{t,x} \Phi_q^k(\pi_q^k)$$

- **Heuristics:**

- Use parameters λ_q , r_q , and \bar{n} to decompose u_q into a “partial wavelet sum,” decompose R_q , Φ_q , π_q by frequency

- **Inductive bounds:**

- Assume bounds for $\|\pi\|_{3/2}$, $\|\pi\|_{\infty}$ (which depend on their frequency)
- All other bounds are *pointwise bounds* in terms of π

- **Scaling law:** Mollifying high-freq. π 's gives rescaled low-freq. π 's

- $\phi_{\lambda^{-1}} * \pi_{\Lambda} = \left(\frac{\lambda}{\Lambda} \right)^{2/3} \pi_{\lambda}, \quad \lambda < \Lambda$

How to propagate spatial support properties (multi-scale pipe flows)

$$\circ u_{q'} = \sum_{q \leq q'} w_q = \sum_{q \leq q'} \sum_k$$

$$\underbrace{a_k^q(t, x)}$$

low-frequency amplitudes,
partition spacetime

$$\underbrace{\mathbb{B}_k^q(\Phi_k^q)}$$

Intermittent Mikado bundle
 $(\partial_t + u_{q-1} \cdot \nabla) \Phi_k^q = 0$

- $\circ \text{supp } a_k^q(t, x) \mathbb{B}_k^q(\Phi_k^q) \cap \text{supp } a_{k'}^q(t, x) \mathbb{B}_{k'}^q(\Phi_{k'}^q) = \emptyset$
- $\circ \text{Either } \text{supp } w_q \cap \text{supp } w_{\tilde{q}} \equiv 0 \text{ or } \text{supp } \hat{w}_q \cap \text{supp } \hat{w}_{\tilde{q}}$

How to propagate spatial support properties (multi-scale pipe flows)

$$\circ u_{q'} = \sum_{q \leq q'} w_q = \sum_{q \leq q'} \sum_k$$

$$\underbrace{a_k^q(t, x)}$$

low-frequency partition
of unity in time and space

$$\underbrace{\mathbb{B}_k^q(\Phi_k^q)}$$

Intermittent Mikado bundle
 $(\partial_t + u_{q-1} \cdot \nabla) \Phi_k^q = 0$

- Intermittent Mikado bundle $\mathbb{B}_k^q = \mathbb{B}_{k,\text{high}}^q \mathbb{B}_{k,\text{low}}^q$ is a multi-scale, intermittent shear flow

How to propagate spatial support properties (multi-scale pipe flows)

$$\circ u_{q'} = \sum_{q \leq q'} w_q = \sum_{q \leq q'} \sum_k$$

$$\underbrace{a_k^q(t, x)}$$

low-frequency partition
of unity in time and space


$$\underbrace{\mathbb{B}_k^q(\Phi_k^q)}$$

Intermittent Mikado bundle
 $(\partial_t + u_{q-1} \cdot \nabla) \Phi_k^q = 0$

- \circ Intermittent Mikado bundle $\mathbb{B}_k^q = \mathbb{B}_{k,\text{high}}^q \mathbb{B}_{k,\text{low}}^q$ is a multi-scale, intermittent shear flow **with flexibility in the support**

How to propagate spatial support properties (multi-scale pipe flows)

- Need to choose the support of $\mathbb{B}_k^q = \mathbb{B}_{k,\text{high}}^q \mathbb{B}_{k,\text{low}}^q$ to dodge any other bundles (for other k' or q') which have overlapping frequency support



Oscillation error and the choice of r_{q+1}

- Simple example, ignoring local energy inequality:

$$R_q^{q-\bar{n}+1} - \pi_q^{q-\bar{n}+1} \text{Id} \rightarrow \begin{pmatrix} R - \pi & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad w_{q+1,R} = \vec{e}_1 (-R + \pi)^{1/2} \underbrace{\mathbb{B}_{q+1,R}(x_2, x_3)}_{\substack{\text{freq's} \\ [\lambda_{q+1} r_{q+1}, \lambda_{q+1}]}}$$

Oscillation error and the choice of r_{q+1}

- Simple example, ignoring local energy inequality:

$$R_q^{q-\bar{n}+1} - \pi_q^{q-\bar{n}+1} \text{Id} \rightarrow \begin{pmatrix} R - \pi & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad w_{q+1,R} = \vec{e}_1 (-R + \pi)^{1/2} \underbrace{\mathbb{B}_{q+1,R}(x_2, x_3)}_{\substack{\text{freq's} \\ [\lambda_{q+1} r_{q+1}, \lambda_{q+1}]}}$$

- Set $\int_{\mathbb{T}^3} \mathbb{B}_{q+1,R}^2 = 1$, use stationarity of $\vec{e}_1 \mathbb{B}_{q+1,R}$, and rewrite

$$\begin{aligned} \text{div}((R - \pi)(\mathbf{e}_1 \otimes \mathbf{e}_1) + w_{q+1,R} \otimes w_{q+1,R}) &= \vec{e}_1 \partial_x \left((R - \pi)(1 - \mathbb{B}_{q+1,R}^2) \right) \\ &= \vec{e}_1 \partial_x (R - \pi) \left(\int_{\mathbb{T}^3} -\text{Id} \right) (\mathbb{B}_{q+1,R}^2) \end{aligned}$$

Oscillation error and the choice of r_{q+1}

- Simple example, ignoring local energy inequality:

$$R_q^{q-\bar{n}+1} - \pi_q^{q-\bar{n}+1} \text{Id} \rightarrow \begin{pmatrix} R - \pi & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad w_{q+1,R} = \vec{e}_1 (-R + \pi)^{1/2} \underbrace{\mathbb{B}_{q+1,R}(x_2, x_3)}_{\substack{\text{freq's} \\ [\lambda_{q+1} r_{q+1}, \lambda_{q+1}]}}$$

- Set $\int_{\mathbb{T}^3} \mathbb{B}_{q+1,R}^2 = 1$, use stationarity of $\vec{e}_1 \mathbb{B}_{q+1,R}$, and rewrite

$$\begin{aligned} \text{div}((R - \pi)(e_1 \otimes e_1) + w_{q+1,R} \otimes w_{q+1,R}) &= \vec{e}_1 \partial_x \left((R - \pi)(1 - \mathbb{B}_{q+1,R}^2) \right) \\ &= \vec{e}_1 \partial_x (R - \pi) \left(\int_{\mathbb{T}^3} -\text{Id} \right) (\mathbb{B}_{q+1,R}^2) \end{aligned}$$

- Put this vector field in divergence form and estimate

Oscillation error and the choice of r_{q+1}

- Simple example:

$$R_q^{q-\bar{n}+1} - \pi_q^{q-\bar{n}+1} \text{Id} \rightarrow \begin{pmatrix} R - \pi & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad w_{q+1,R} = \vec{e}_1 (-R + \pi)^{1/2} \underbrace{\mathbb{B}_{q+1,R}(x_2, x_3)}_{\substack{\text{freq's} \\ [\lambda_{q+1} r_{q+1}, \lambda_{q+1}]}}$$

- Estimate error in $L^{3/2}$ (since R, π scale like velocity squared)

$$\left\| \text{div}^{-1} \left(\vec{e}_1 \partial_x (R - \pi) \left(\text{Id} - \int_{\mathbb{T}^3} \right) (\mathbb{B}_{q+1,R}^2) \right) \right\|_{3/2}$$

Oscillation error and the choice of r_{q+1}

- Simple example:

$$R_q^{q-\bar{n}+1} - \pi_q^{q-\bar{n}+1} \text{Id} \rightarrow \begin{pmatrix} R - \pi & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad w_{q+1,R} = \vec{e}_1 (-R + \pi)^{1/2} \underbrace{\mathbb{B}_{q+1,R}(x_2, x_3)}_{\substack{\text{freq's} \\ [\lambda_{q+1} r_{q+1}, \lambda_{q+1}]}}$$

- Estimate error in $L^{3/2}$ (since R, π scale like velocity squared)

$$\left\| \text{div}^{-1} \left(\vec{e}_1 \partial_x (R - \pi) \left(\text{Id} - \int_{\mathbb{T}^3} \right) (\mathbb{B}_{q+1,R}^2) \right) \right\|_{3/2}$$

- Requires knowledge of $\|\mathbb{B}_{q+1,R}\|_3$, which *grows* as $r_{q+1} \rightarrow 0$ from Bernstein's inequality and $\int \mathbb{B}_{q+1,R}^2 = 1$

Oscillation error and the choice of r_{q+1}

- Simple example:

$$R_q^{q-\bar{n}+1} - \pi_q^{q-\bar{n}+1} \text{Id} \rightarrow \begin{pmatrix} R - \pi & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad w_{q+1,R} = \vec{e}_1 (-R + \pi)^{1/2} \underbrace{\mathbb{B}_{q+1,R}(x_2, x_3)}_{\substack{\text{freq's} \\ [\lambda_{q+1} r_{q+1}, \lambda_{q+1}]}}$$

- Estimate in $L^{3/2}$ (since R, π scale like velocity squared)

$$\left\| \text{div}^{-1} \left(\vec{e}_1 \partial_x (R - \pi) \left(\text{Id} - \int_{\mathbb{T}^3} \right) (\mathbb{B}_{q+1,R}^2) \right) \right\|_{3/2}$$

- Requires knowledge of $\|\mathbb{B}_{q+1,R}\|_3$, which *grows* as $r_{q+1} \rightarrow 0$ from Bernstein's inequality and $\int \mathbb{B}_{q+1,R}^2 = 1$
- Conclusion:** Nonlinear error prefers $r_{q+1} = 1$, i.e. *less* intermittency

Scaling law: $\pi_\Lambda * \phi_{\lambda^{-1}} = \left(\frac{\lambda}{\Lambda}\right)^{2/3} \pi_\lambda, \lambda < \Lambda$

- Simple example (requires $R \leq \pi$):

$$R_q^{q-\bar{n}+1} - \pi_q^{q-\bar{n}+1} \text{Id} \rightarrow \begin{pmatrix} R - \pi & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad w_{q+1,R} = \vec{e}_1 (-R + \pi)^{1/2} \underbrace{\mathbb{B}_{q+1,R}(x_2, x_3)}_{\substack{\text{freq's} \\ [\lambda_{q+1} r_{q+1}, \lambda_{q+1}]}}$$

Scaling law: $\pi_\Lambda * \phi_{\lambda^{-1}} = \left(\frac{\lambda}{\Lambda}\right)^{2/3} \pi_\lambda, \lambda < \Lambda$

- Simple example (requires $R \leq \pi$):

$$R_q^{q-\bar{n}+1} - \pi_q^{q-\bar{n}+1} \text{Id} \rightarrow \begin{pmatrix} R - \pi & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad w_{q+1,R} = \vec{e}_1 (-R + \pi)^{1/2} \underbrace{\mathbb{B}_{q+1,R}(x_2, x_3)}_{\substack{\text{freq's} \\ [\lambda_{q+1} r_{q+1}, \lambda_{q+1}]}}$$

- We need to build π_Λ such that

$$\pi_\Lambda \geq \left| \text{div}^{-1} \left(\vec{e}_1 \partial_x (R - \pi) \left(\text{Id} - \underset{\mathbb{T}^3}{f} \right) (\mathbb{B}_{q+1,R}^2) \right) \right|$$

Scaling law: $\pi_\Lambda * \phi_{\lambda^{-1}} = \left(\frac{\lambda}{\Lambda}\right)^{2/3} \pi_\lambda, \lambda < \Lambda$

- Simple example (requires $R \leq \pi!!$):

$$R_q^{q-\bar{n}+1} - \pi_q^{q-\bar{n}+1} \text{Id} \rightarrow \begin{pmatrix} R - \pi & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad w_{q+1,R} = \vec{e}_1 (-R + \pi)^{1/2} \underbrace{\mathbb{B}_{q+1,R}(x_2, x_3)}_{\substack{\text{freq's} \\ [\lambda_{q+1} r_{q+1}, \lambda_{q+1}]}}$$

- We need to build π_Λ such that

$$\pi_\Lambda \geq \left| \text{div}^{-1} \left(\vec{e}_1 \partial_x (R - \pi) \left(\text{Id} - f_{\mathbb{T}^3} \right) (\mathbb{B}_{q+1,R}^2) \right) \right|$$

- Assuming div^{-1} gains a power of high frequency $\Lambda \approx \lambda_{q+1} r_{q+1}$, and using the inductive assumption $|\nabla_x R| + |\nabla_x \pi| \lesssim \lambda \pi$, we set

$$\begin{aligned} \pi_\Lambda &= \frac{\pi \lambda}{\Lambda} \left[f + \left(\text{Id} - f \right) \right] \left| \left(\text{Id} - f \right) (\mathbb{B}_{q+1,R}^2) \right| \\ &= \frac{\pi \lambda}{\Lambda} + \text{high frequency} \end{aligned}$$

Linear errors and the choice of r_{q+1}

- Recall that

$$w_{q+1,R} = \vec{e}_1(-R + \pi)^{1/2} \underbrace{\mathbb{B}_{q+1,R}(x_2, x_3)}_{\substack{\text{freq's} \\ [\lambda_{q+1}r_{q+1}, \lambda_{q+1}]}}$$

and consider the (linear) Nash error

$$R_{q+1}^{\text{Nash}} = \text{div}^{-1}(w_{q+1,R} \cdot \nabla u_q) \quad \implies \quad w_{q+1,R} \cdot \nabla u_q = \text{div} \left(R_{q+1}^{\text{Nash}} \right)$$

Linear errors and the choice of r_{q+1}

- Recall that

$$w_{q+1,R} = \vec{e}_1(-R + \pi)^{1/2} \underbrace{\mathbb{B}_{q+1,R}(x_2, x_3)}_{\substack{\text{freq's} \\ [\lambda_{q+1} r_{q+1}, \lambda_{q+1}]}}$$

and consider the (linear) Nash error

$$R_{q+1}^{\text{Nash}} = \operatorname{div}^{-1}(w_{q+1,R} \cdot \nabla u_q) \quad \implies \quad w_{q+1,R} \cdot \nabla u_q = \operatorname{div} \left(R_{q+1}^{\text{Nash}} \right)$$

- Since it is linear, and $\mathbb{B}_{q+1,R}$ is L^2 -normalized it prefers *more* intermittency, so that

$$\|\mathbb{B}_{q+1,R}\|_{L^{3/2}} \rightarrow 0 \quad \text{as} \quad r_{q+1} \rightarrow 0$$

Linear errors and the choice of r_{q+1}

- Recall that

$$w_{q+1,R} = \vec{e}_1(-R + \pi)^{1/2} \underbrace{\mathbb{B}_{q+1,R}(x_2, x_3)}_{\substack{\text{freq's} \\ [\lambda_{q+1} r_{q+1}, \lambda_{q+1}]}}$$

and consider the (linear) Nash error

$$R_{q+1}^{\text{Nash}} = \text{div}^{-1}(w_{q+1,R} \cdot \nabla u_q) \quad \implies \quad w_{q+1,R} \cdot \nabla u_q = \text{div} \left(R_{q+1}^{\text{Nash}} \right)$$

- Since it is linear, and $\mathbb{B}_{q+1,R}$ is L^2 -normalized it prefers *more* intermittency, so that

$$\|\mathbb{B}_{q+1,R}\|_{L^{3/2}} \rightarrow 0 \quad \text{as} \quad r_{q+1} \rightarrow 0$$

- Recall that the nonlinear error preferred *less* intermittency, i.e. $r_{q+1} = 1$

Linear errors and the choice of r_{q+1}

- Recall that

$$w_{q+1,R} = \vec{e}_1(-R + \pi)^{1/2} \underbrace{\mathbb{B}_{q+1,R}(x_2, x_3)}_{\substack{\text{freq's} \\ [\lambda_{q+1} r_{q+1}, \lambda_{q+1}]}}$$

and consider the (linear) Nash error

$$R_{q+1}^{\text{Nash}} = \text{div}^{-1}(w_{q+1,R} \cdot \nabla u_q) \quad \implies \quad w_{q+1,R} \cdot \nabla u_q = \text{div} \left(R_{q+1}^{\text{Nash}} \right)$$

- Since it is linear, and $\mathbb{B}_{q+1,R}$ is L^2 -normalized it prefers *more* intermittency, so that

$$\|\mathbb{B}_{q+1,R}\|_{L^{3/2}} \rightarrow 0 \quad \text{as} \quad r_{q+1} \rightarrow 0$$

- Recall that the nonlinear error preferred *less* intermittency, i.e. $r_{q+1} = 1$
- Heuristic estimates of this choice dictate a single acceptable choice

$$r_{q+1}^{\text{Goldilocks}} = \left(\frac{\text{freq. of } R}{\text{max. freq. of } \mathbb{B}_{q+1,R}} \right)^{1/2}$$

Open problems

- (Dream Theorem) Construct a sequence of suitable solutions to Navier-Stokes converging to a dissipative Euler solution in $B_{3,\infty}^{1/3-}$

Open problems

- (Dream Theorem) Construct a sequence of suitable solutions to Navier-Stokes converging to a dissipative Euler solution in $B_{3,\infty}^{1/3-}$
- (Partial progress) Construct solutions to Navier-Stokes in H^β for β non-negligible

Open problems

- (Dream Theorem) Construct a sequence of suitable solutions to Navier-Stokes converging to a dissipative Euler solution in $B_{3,\infty}^{1/3-}$
- (Partial progress) Construct solutions to Navier-Stokes in H^β for β non-negligible
- Construct wild solutions to 2D Euler with Lebesgue integrable vorticity (preferably for p as large as possible!)

Open problems

- (Dream Theorem) Construct a sequence of suitable solutions to Navier-Stokes converging to a dissipative Euler solution in $B_{3,\infty}^{1/3-}$
- (Partial progress) Construct solutions to Navier-Stokes in H^β for β non-negligible
- Construct wild solutions to 2D Euler with Lebesgue integrable vorticity (preferably for p as large as possible!)
- Construct wild solutions to 3D Euler with well-defined helicity $\int_{\mathbb{T}^3} u \cdot (\nabla \times u)$

Thanks for your attention!