

## Homework 5 – SPERNER’S THEOREM AND UNIT DISTANCES

Janos Pach &amp; Nabil Mustafa

24 March

## Questions

1. Given a set system  $\mathcal{F}$  over the base set  $\{1, \dots, n\}$ , we call  $\mathcal{F}$  *semi-independent* if it contains no three sets  $A, B, C$  such that  $A \subset B \subset C$ . Prove that  $|\mathcal{F}| \leq 2^{\lfloor n/2 \rfloor}$ .
2. Let  $a_1, \dots, a_n$  be real numbers with  $|a_i| \geq 1$ . Let  $p(a_1, \dots, a_n)$  be the number of vectors  $(\epsilon_1, \dots, \epsilon_n)$ , where  $\epsilon_i = \pm 1$ , such that

$$-1 < \sum_{i=1}^n \epsilon_i a_i < 1$$

Prove that for any  $a_1, \dots, a_n$ , we have  $p(a_1, \dots, a_n) \leq 2^{\lfloor n/2 \rfloor}$ .

3. Let  $X$  be an  $n$ -element set, and let  $S_1, \dots, S_n$  be subsets of  $X$  such that  $|S_i \cap S_j| \leq 1$  for all  $1 \leq i < j \leq n$ . Prove that at least one set has size at most  $C\sqrt{n}$  for some absolute constant  $C$ .
4. Let  $t(j)$  denote the number of divisors of the number  $j$ . Give an expression for the number  $\sum_{i=1}^n t(i)$ .
5. Given a set  $P$  of  $n$  points, and a set  $L$  of  $n$  lines in the plane, an incidence is a pair  $(p, l)$ , where  $p \in P$ ,  $l \in L$ , and the point  $p$  lies on the line  $l$ . Prove that given *any* set of  $n$  distinct lines  $L$  and  $n$  distinct points  $P$ , the number of incidences are at most  $3n^{1.5}$ .

**Bonus Problem.** You are given a set  $P$  of 10 integers from the set  $\{1, \dots, 100\}$ . Prove that one can always find two *disjoint* subsets of  $P$  such that the sum of the elements in the two sets are equal. For example, given the set  $\{8, 15, 23, 59, 61, 70, 75, 88, 91, 97\}$ , the two sets are  $\{8, 15, 97\}$  and  $\{59, 61\}$ .

10 points.