

Last name:	First name:																																				
Exercise:	<table border="1" style="border-collapse: collapse; width: 100%;"> <tr> <td style="width: 12.5%;">1</td> <td style="width: 12.5%;">2</td> <td style="width: 12.5%;">3</td> <td style="width: 12.5%;">4</td> <td style="width: 12.5%;">5</td> <td style="width: 12.5%;">6</td> <td style="width: 12.5%;">7</td> <td style="width: 12.5%;">8</td> <td style="width: 12.5%;">Σ</td> </tr> <tr> <td style="text-align: center;">10</td> <td style="text-align: center;">10</td> <td style="text-align: center;">10</td> <td style="text-align: center;">10</td> <td style="text-align: center;">10</td> <td style="text-align: center;">10</td> <td style="text-align: center;">10</td> <td style="text-align: center;">10</td> <td style="text-align: center;">50</td> </tr> <tr> <td style="text-align: center;"> </td> <td style="text-align: center;"> </td> <td style="text-align: center;"> </td> <td style="text-align: center;"> </td> <td style="text-align: center;"> </td> <td style="text-align: center;"> </td> <td style="text-align: center;"> </td> <td style="text-align: center;"> </td> <td style="text-align: center;"> </td> </tr> <tr> <td style="text-align: center;"> </td> <td style="text-align: center;"> </td> <td style="text-align: center;"> </td> <td style="text-align: center;"> </td> <td style="text-align: center;"> </td> <td style="text-align: center;"> </td> <td style="text-align: center;"> </td> <td style="text-align: center;"> </td> <td style="text-align: center;"> </td> </tr> </table>	1	2	3	4	5	6	7	8	Σ	10	10	10	10	10	10	10	10	50																		
1	2	3	4	5	6	7	8	Σ																													
10	10	10	10	10	10	10	10	50																													
max points:																																					
achieved points:																																					
chosen exercises:																																					

Check whether the exam is complete: It should have 12 pages (Exercises 1–8). Write your name on the title page. Solutions have to be written below the exercises. Solutions must be comprehensible. In case of lack of space, you can ask for additional paper from the exam supervision. Please put your name on each additional sheet and indicate which exercise it belongs to.

Use neither pencil nor red colored pen!

Duration: 180 min

Grading:

Every exercise gives 10 points, and you are supposed to solve 5 of them. There are 6 exercises marked with [*] and two exercises marked with [Δ]. Math students can choose among the [*]-exercises. Non-math students can choose among all exercises. **Please mark the 5 exercises you have chosen in the tabular above!**

You are allowed to bring a pocket calculator and an A4-“cheat-sheet”.

Exercise 1 [*]: (Multiple Choice, points $\{-1, 0, 1\}$ each)

No justifications needed. Mark 'yes' or 'no'. **Wrong answers cause negative points!**

<p>a) Given a set $A := \{a_1, \dots, a_m\} \subset \mathbb{R}^n$, then for all $x \in \text{conv}(A)$ there are $\lambda_1, \dots, \lambda_m \in \mathbb{R}_{\geq 0}$, at most n of them nonzero, such that $\sum_{i=1}^m \lambda_i = 1$ and $x = \sum_{i=1}^m \lambda_i a_i$.</p>	<p><input type="radio"/> yes <input type="radio"/> no</p>
<p>b) Given a linear program $\max\{c^T x \mid Ax \leq b\}$, with $A \in \mathbb{R}^{m \times n}, b \in \mathbb{R}^m, c \in \mathbb{R}^n$ and $m > n$. If x is an optimal solution, at least n of the LPs inequalities are satisfied with equality by x.</p>	<p><input type="radio"/> yes <input type="radio"/> no</p>
<p>c) Given a linear program $\max\{c^T x \mid Ax \leq b\}$, with $A \in \mathbb{R}^{m \times n}, b \in \mathbb{R}^m, c \in \mathbb{R}^n$. If the LP is feasible and bounded, then there is a <i>roof</i> B such that its vertex is an optimal solution of the LP.</p>	<p><input type="radio"/> yes <input type="radio"/> no</p>
<p>d) For a linear program $\max\{c^T x \mid Ax \leq b\}$, with $A \in \mathbb{R}^{m \times n}, b \in \mathbb{R}^m, c \in \mathbb{R}^n$, let j be a row that entered the roof in the ith iteration of the simplex algorithm. Then it cannot leave the roof in the $(i + 1)$st iteration.</p>	<p><input type="radio"/> yes <input type="radio"/> no</p>
<p>e) For a linear program $\max\{c^T x \mid Ax \leq b\}$, with $A \in \mathbb{R}^{m \times n}, b \in \mathbb{R}^m, c \in \mathbb{R}^n$, let j be a row that has left the roof in the ith iteration of the simplex algorithm. Then it cannot reenter into the roof in the $(i + 1)$st iteration.</p>	<p><input type="radio"/> yes <input type="radio"/> no</p>
<p>f) Given a linear program $\max\{c^T x \mid Ax \leq b\}$, with $A \in \mathbb{Z}^{m \times n}, b \in \mathbb{Z}^m, c \in \mathbb{R}^n$, A full column rank. If A is totally unimodular, then the vertex of every roof of the LP is an integer vector.</p>	<p><input type="radio"/> yes <input type="radio"/> no</p>
<p>g) Consider an integer program $\max\{c^T x \mid Ax \leq b, x \in \mathbb{Z}^n\}$, with $A \in \mathbb{Z}^{m \times n}, b \in \mathbb{Z}^m, c \in \mathbb{R}^n$. Given an optimal solution x^* to the IP, there is a solution y^* to the IP</p> $\min\{b^T y \mid A^T y = c, y \in \mathbb{Z}_{\geq 0}^m\}$ <p>such that $c^T x^* = b^T y^*$.</p>	<p><input type="radio"/> yes <input type="radio"/> no</p>
<p>h) There is a linear program</p> $\max\{c^T x : Ax \leq b\},$ <p>with $A \in \mathbb{R}^{m \times n}, b \in \mathbb{R}^m, c \in \mathbb{R}^n$ such that both the LP and its dual are infeasible.</p>	<p><input type="radio"/> yes <input type="radio"/> no</p>
<p>i) Given a directed graph $G = (V, A)$ and a length function $\ell : A \rightarrow \mathbb{Z}$. One can decide if a negative cycle in G (w.r.t. length ℓ) exists in time $O(V \cdot A)$.</p>	<p><input type="radio"/> yes <input type="radio"/> no</p>
<p>j) Given a directed graph $G = (V, A)$ and two nodes $s, t \in V$. The maximum number of arc-disjoint $s - t$-paths is equal to the smallest number of arcs in an $s - t$-cut. [A collection P_1, \dots, P_k is arc-disjoint, if no pair of paths has an arc in common.]</p>	<p><input type="radio"/> yes <input type="radio"/> no</p>

Exercise 2 [Δ] (LP modelling 1):

A refinery can produce gas of two types of qualities A and B, composed of three kinds of crude oil 1, 2, 3. To produce one unit of gas A or B, one can combine fractional quantities of the three kinds of crude oils such that they sum up to 1. The following restrictions apply: At most 30% of crude oil 1, at least 40% of crude oil 2 and at most 50% of crude oil 3 has to be used to produce gas A. At most 50% of crude oil 1 and at least 10% of crude oil 2 has to be used to produce gas B (There are no restrictions for crude oil 3 here). The total quantity of each crude oil that can be used in production is limited, and it comes with a cost per unit which is as follows:

crude oil	max quantity	cost per unit
1	3000	3
2	2000	6
3	4000	4

One unit of gas A sells for 5.5, and one unit of gas B sells for 4.5.

Formulate the problem of maximizing revenue (income by selling the gas minus cost of raw material) as a linear program.

You do not need to transform you LP into standard form!

Solution:

Use reverse side if you need more space

Exercise 3 [*] (LP modelling 2):

Consider the polyhedron

$$P := \{x \in \mathbb{R}^n : Ax \leq b\}$$

for some $b \in \mathbb{R}^m$, $A \in \mathbb{R}^{m \times n}$.

A ball with center y and radius r is defined as the set:

$$B_r(y) := \{x \in \mathbb{R}^n : \|x - y\|_2 \leq r\}.$$

We want to find a ball (i.e. its center y) of largest possible radius which is entirely contained in P .

Formulate this problem as a linear program. **Prove that your LP formulation gives the desired result!**

Hints:

- For $v \in \mathbb{R}^n$, $\|v\|_2^2 = v^T v$.
- Consider each row of $a_i x \leq b_i$ individually. Find a necessary and sufficient condition that asserts that $B_r(y)$ does not violate $a_i x \leq b_i$.

Solution:

Use reverse side if you need more space

Exercise 4 [*] (LP Duality / Complementary slackness):

(a) Consider a linear program

$$\max\{c^T x : Ax \leq b\}$$

for some $c \in \mathbb{R}^n$, $b \in \mathbb{R}^m$, $A \in \mathbb{R}^{m \times n}$.

Prove the complementary slackness theorem: Given an optimal solution x^* for the LP, there is an optimal solution y^* for its dual such that $y_i^* = 0$ for all rows i of the primal that are not satisfied with equality by x^* .

Now consider the following LP:

$$\begin{array}{rcll} \max & 10x_1 & + & 4x_2 \\ & x_1 & & \leq -4 \\ & & & x_2 \leq -6 \\ & x_1 & + & 2x_2 \leq -20 \\ & 2x_1 & + & x_2 \leq -17 \end{array}$$

(b) Write down a dual of the LP in standard form.

(c) Show that $x^* := (-4, -9)^T$ is an optimal solution for the LP by giving a suitable solution for the dual LP (using the complementary slackness theorem).

Solution:

Use the next page if you need more space

Solution:

Use reverse side if you need more space

Exercise 5 [*] (Roofs):

Consider a linear program

$$\max\{c^T x : Ax \leq b\}$$

for some $c \in \mathbb{R}^n$, $b \in \mathbb{R}^m$, $A \in \mathbb{R}^{m \times n}$ where A has full column rank.

Consider a subset $B \subseteq \{1, \dots, m\}$ of rows such that $|B| = n$ and the rows a_i , $i \in B$ are linearly independent.

Show that if $c \in \text{cone}\{a_i : i \in B\}$, then B is a roof.

Solution:

Use reverse side if you need more space

Solution:

Use reverse side if you need more space

Exercise 7 [*] (Flow modeling):

There are k projects available to be undertaken. Let P be the set of projects. Each project $i \in P$ has associated a positive revenue $r_i \in \mathbb{N}$ and requires a set $S_i \subseteq R$ of resources to be available. Each resource $j \in R$ has an associated cost $c_j \in \mathbb{N}$.

If a resource $j \in R$ is required by several projects, it needs to be bought only once.

Show how to use a max-flow (min-cut) algorithm to solve the problem of choosing a subset of projects, such that the sum of the revenues minus the cost of the required resources is maximized.


Hint: Construct a graph such that every cut of finite capacity corresponds to a set of selected projects and all their required resources. (In particular, it might be useful to put a capacity of ∞ on some arcs.)

Solution:

Use reverse side if you need more space

Exercise 8 [Δ] (Max $s-t$ -flows):

Consider the following graph $G = (V, A)$. The labels on the arcs $a \in A$ are of the form $f(a)/u(a)$, i.e. they define functions $f : A \rightarrow \mathbb{Q}_{\geq 0}$ and $u : A \rightarrow \mathbb{Q}_{\geq 0}$.



- Argue why f is a *feasible* $s-t$ -flow in G subject to the capacities u . What is the value of the flow?
- Perform the Ford-Fulkerson algorithm to compute a maximum $s-t$ -flow in G . For each iteration give the residual network. You can start with the flow f . Give the flow, its value and a minimum $s-t$ cut.
- Prove the weak duality theorem: Given a feasible $s-t$ -flow f and an $s-t$ -cut $\delta^{out}(U)$ for some $U \subset V$, then

$$\text{value}(f) \leq u(\delta^{out}(U)).$$

Hint: You can use the fact that $\text{excess}_f(U) = \sum_{v \in U} \text{excess}_f(v)$ for each $U \subseteq V$.

Solution:

Use the next page if you need more space

Solution:

Use reverse side if you need more space