

Neural Networks and Biological Modeling

Professor Wulfram Gerstner
Laboratory of Computational Neuroscience

QUESTION SET 8

Exercise 1: Poisson neuron

We consider a neuron that fires stochastically. Its firing rate is described by a Poisson process of rate ρ . In other words, in every small time interval Δt , the probability that the neuron fires is given by $\rho\Delta t$.

1.1 What is the probability that the neuron does *not* fire during a time of arbitrarily large length t ?
Hint: Consider first the probability of not firing during a short interval Δt .

1.2 Suppose that the neuron has fired at time t_0 . Calculate the distribution of intervals $P(s)$, i.e., the probability density that the neuron fires its next spike after a time s .

1.3 Suppose that the neuron is driven by some input. For $t < t_0$, the input is weak, so that its firing rate is $\rho_0 = 2\text{Hz}$. For $t_0 < t < t_1 = t_0 + 100\text{ms}$, the input is strong and the neuron fires at $\rho_1 = 20\text{Hz}$.

(i) Calculate the interval distributions for weak and strong stimuli.

(ii) What is the probability of having a “burst” consisting of two intervals of less than 20 ms each if the input is weak/strong?

(iii) Suppose that the onset time t_0 of the strong input is unknown; can an observer, who is looking at the neuron’s output, decide whether the input is weak or strong?

1.4 Suppose that a Poisson neuron with a constant rate of 20 Hz emits in a trial of 5 second duration 100 spikes at times $t^{(1)}, t^{(2)}, \dots, t^{(100)}$. The experiment is repeated such that a second spike train with a duration of 5 seconds is observed.

What is the percentage of spikes that coincide between the first and second trial with a precision of $\pm 2\text{ms}$? More generally, what percentage of spikes coincide between two trials of a Poisson neuron with arbitrary rate ρ_0 under the assumption that trials are sufficiently long?

Exercise 2: Stochastic spike arrival

Consider a neuron with a passive membrane,

$$\tau \frac{du}{dt} = -(u - u_{\text{rest}}) + RI(t). \quad (1)$$

2.1 The neuron receives synaptic input at a rate ν such that

$$I(t) = q \sum_f \delta(t - t^f). \quad (2)$$

Calculate the average value of membrane potential as a function of the presynaptic rate ν , assuming

stochastic (Poisson) spike arrival.

Hint: Integrate Eq. 1 keeping explicitly the δ -function. Under the assumption of stochastic spike arrival we have $\left\langle \sum_f \delta(t - t^f) \right\rangle = \nu$.

2.2 Calculate the average value of membrane potential as a function of the presynaptic rate ν if the current coming from the presynaptic activity is:

$$I(t) = \sum_f \alpha(t - t^f). \quad (3)$$

Hint: As before, integrate Eq. 1 keeping the δ -function explicit.

Exercise 3: Renewal process

We consider a neuron with relative refractoriness. Given an output spike at time \hat{t} , the probability of firing is given by

$$\rho(t - \hat{t}) = \begin{cases} 0 & \text{for } t - \hat{t} < t_{\text{abs}} \\ [t - \hat{t} - t_{\text{abs}}] \frac{\rho_0}{2} & \text{for } t_{\text{abs}} < t - \hat{t} < t_{\text{abs}} + 2 \\ \rho_0 & \text{otherwise.} \end{cases} \quad (4)$$

Calculate the survivor function and the interval distribution.

Exercise 4: Homework

4.1 The poisson neuron has a probability to fire in a very small interval Δt equal to $\nu \Delta t$. What will be the probability to observe exactly k spikes in the time interval $T = N \Delta t$ ($P_k(T)$)? Start with the probability to observe k events in N slots (the binomial distribution):

$$P(k, N) = \frac{N!}{k!(N - k)!} p_1^{N-k} p_2^k$$

where p_1 and p_2 are the probability to spike or to remain silent in one Δt slot. Take the continuous time limit with Stirling's approximation ($N! \approx (N/e)^N$ for large N) to obtain the Poisson distribution:

$$P_N = \frac{(\nu T)^N}{N!} \exp(-\nu T)$$

Verify that this distribution predicts an average number of spikes $\langle N \rangle = \nu T$.