ÉCOLE POLYTECHNIQUE FÉDÉRALE DE LAUSANNE
School of Computer and Communication Sciences

Handout 7 Information Theory and Coding
Solutions to Homework 3 Oct. 11, 2021

PROBLEM 1.

(a) Recall that $\mathcal{C}$ is uniquely decodable means that $\mathcal{C}^*$ is injective, i.e., for any $u^n \neq v^m$ we have $\mathcal{C}^n(u^n) \neq \mathcal{C}^m(v^m)$. In particular, whenever $u^n \neq v^n$ we have $\mathcal{C}^n(u^n) \neq \mathcal{C}^n(v^n)$. The last statement is the definition of $\mathcal{C}^n$ being injective.

(b) Since we are supposed to show that $u_1 \neq v_1$, we may assume that $|\mathcal{U}| \geq 2$.

If $\mathcal{C}$ is not uniquely decodable, then there are $u^n \neq v^m$ such that $\mathcal{C}^n(u^n) = \mathcal{C}^m(v^m)$. Among all such (u^n, v^m) choose one for which $n+m$ is smallest, and assume (without loss of generality) that $m \leq n$. If $m \geq 1$ we are done, since in this case we must have $u_1 \neq v_1$ (because, if not, we can replace u^n by $\tilde{u}^{n-1} = u_2 \dots u_n$ and v^m by $\tilde{v}^{m-1} = v_2 \dots v_m$, contradicting $m+n$ being smallest).

Otherwise, $m = 0$ and $v^m = \lambda$ (the null string) with $\mathcal{C}(v^m) = \lambda$. Since $u^n \neq v^m = \lambda$ and $\mathcal{C}(u^n) = \lambda$, we have a letter $a = u_1 \in \mathcal{U}$ such that $\mathcal{C}(a) = \lambda$. Take now any letter $b \in \mathcal{U}$ with $b \neq a$, and note that $\mathcal{C}^2(ab) = \mathcal{C}^1(b)$, i.e., there are two source sequences that differ in their first letter and have the same representation.

(c) $\mathcal{C}$ is not uniquely decodable means that there is $u^n \neq v^m$ such that $\mathcal{C}^n(u^n) = \mathcal{C}^m(v^m)$. If $n = m$ then we are done: this would by definition mean that be $\mathcal{C}^n$ is not injective. If $n \neq m$, we could attempt the following reasoning: observe $\mathcal{C}^*(u^n v^m) = \mathcal{C}^*(v^m u^n)$ and conclude that $\mathcal{C}^{m+n}$ is not injective. However this reasoning fails because we can't be sure that $u^n v^m \neq v^m u^n$ just because $u^n \neq v^m$. (E.g., suppose $u^n = a$ and $v^m = aa$). This is the reason the problem has "part (b)":

As $\mathcal{C}$ is not uniquely decodable, we can find u^n and v^m as in part (b). Now observe that (i) $u^n v^m \neq v^m u^n$ (as they differ in their first letter), (ii) $u^n v^m$ and $v^m u^n$ have the same length $k = n + m$, and $\mathcal{C}^k(u^n v^m) = \mathcal{C}^k(v^m u^n)$, i.e., $\mathcal{C}^k$ is not singular.

Moral of the problem: it is clear that the statement "$\mathcal{C}^*$ is injective" is a stronger statement than "for every n, $\mathcal{C}^n$ is injective" — since the first ensures that $u^n \neq v^m$ are assigned different codewords not only when $n = m$ but also for $n \neq m$ — so part (a) is unsurprising. The statement "$\mathcal{C}^n$ is injective for each n" only means that different source sequences of *same length* get different representations; it is not immediately clear that this will also imply that source sequences of *different lengths* also get different representations. Part (c) shows this is indeed the case: that injectiveness of $\mathcal{C}^n$ for every n implies the injectiveness of $\mathcal{C}^*$.

Problem 2.

(a) We already know that

$$H(X) + H(Y) \geq H(XY),$$
$$H(Y) + H(Z) \geq H(YZ),$$

and

$$H(Z) + H(X) \geq H(ZX).$$

Adding these inequalities together and diving by two gives

$$H(X) + H(Y) + H(Z) \geq \frac{1}{2}\big[H(XY) + H(YZ) + H(ZX)\big].$$

(b) The difference between the left and right sides, i.e.,

$$H(XY) + H(YZ) - H(XYZ) - H(Y),$$

equals

$$H(X|Y) - H(X|YZ) = I(X; Z|Y),$$

which is always positive.

(c) Using (b) with (YZX) and (ZXY) in the role of (XYZ) gives the inequalities

$$H(YZ) + H(ZX) \geq H(XYZ) + H(Z)$$

and

$$H(ZX) + H(XY) \geq H(XYZ) + H(X).$$

Adding the inequality in (b) to these two gives

$$2\big[H(XY) + H(YZ) + H(ZX)\big] \geq 3H(XYZ) + H(X) + H(Y) + H(Z).$$

(d) Since $H(X) + H(Y) + H(Z) \geq H(XYZ)$, (c) yields

$$2\big[H(XY) + H(YZ) + H(ZX)\big] \geq 4H(XYZ).$$

(e) Let $\big\{(x_i, y_i, z_i) : i = 1, \ldots, n\big\}$ be the xyz-coordinates of the n points. Let X, Y and Z be random variables with $\Pr\big((X, Y, Z) = (x_i, y_i, z_i)\big) = 1/n$ for every $1 \leq i \leq n$. Then, $H(XYZ) = \log_2 n$. Furthermore, the random pair (XY) takes values in the projection of the n points to the xy plane and similarly for (YZ) and (ZX). Thus $H(XY) \leq \log_2 n_{xy}$, $H(YZ) \leq \log_2 n_{yz}$, and $H(ZX) \leq \log_2 n_{zx}$. Part (d) now yields

$$\log_2[n_{xy} n_{yz} n_{zx}] \geq H(XY) + H(YZ) + H(ZX) \geq 2H(XYZ) = 2\log_2 n,$$

which implies that $n_{xy} n_{yz} n_{zx} \geq n^2$.

The relationship between $H(XYZ)$ and $H(XY)$, $H(YZ)$ and $H(ZX)$ is a special case of Han's inequality, which, for a collection of n random variables relates the sum of the $\binom{n}{k}$ joint entropies of k out of n random variables to the sum of the $\binom{n}{k+1}$ entropies of $k+1$ out of n random variables.

The combinatorial fact about the projections of points in 3D is known as Shearer's lemma.

Problem 3.

$$\begin{aligned} H(X) &= -\sum_{k=1}^{M} P_X(a_k)\log P_X(a_k) \\ &= -\sum_{k=1}^{M-1}(1-\alpha)P_Y(a_k)\log[(1-\alpha)P_Y(a_k)] - \alpha\log\alpha \\ &= (1-\alpha)H(Y) - (1-\alpha)\log(1-\alpha) - \alpha\log\alpha \end{aligned}$$

Since Y is a random variable that takes $M-1$ values $H(Y) \le \log(M-1)$ with equality if and only if Y takes each of its possible values with equal probability.

Problem 4.

(a) Using the chain rule for mutual information,

$$I(X,Y;Z) = I(X;Z) + I(Y;Z \mid X) \ge I(X;Z),$$

with equality iff $I(Y;Z \mid X) = 0$, that is, when Y and Z are conditionally independent given X.

(b) Using the chain rule for conditional entropy,

$$H(X,Y \mid Z) = H(X \mid Z) + H(Y \mid X,Z) \ge H(X \mid Z),$$

with equality iff $H(Y \mid X,Z) = 0$, that is, when Y is a function of X and/or Z.

(c) Using first the chain rule for entropy and then the definition of conditional mutual information,

$$\begin{aligned} H(X,Y,Z) - H(X,Y) = H(Z \mid X,Y) &= H(Z \mid X) - I(Y;Z \mid X) \\ &\le H(Z \mid X) = H(X,Z) - H(X)\,, \end{aligned}$$

with equality iff $I(Y;Z \mid X) = 0$, that is, when Y and Z are conditionally independent given X.

(d) Using the chain rule for mutual information,

$$I(X;Z \mid Y) + I(Z;Y) = I(X,Y;Z) = I(Z;Y \mid X) + I(X;Z)\,,$$

and therefore

$$I(X;Z \mid Y) = I(Z;Y \mid X) - I(Z;Y) + I(X;Z)\,.$$

We see that this inequality is actually an equality in all cases.

Problem 5. Let X^i denote $X_1, \ldots, X_i$.

(a) By stationarity we have for all $1 \le i \le n$,

$$H(X_n|X^{n-1}) \le H(X_n|X_{n-i+1}, X_{n-i+2}, \ldots, X_{n-1}) = H(X_i|X^{i-1}),$$

which implies that,

$$\begin{aligned} H(X_n|X^{n-1}) &= \frac{\sum_{i=1}^{n} H(X_n|X^{n-1})}{n} && (1) \\ &\le \frac{\sum_{i=1}^{n} H(X_i|X^{i-1})}{n} && (2) \\ &= \frac{H(X_1, X_2, \ldots, X_n)}{n}. && (3) \end{aligned}$$

(b) By the chain rule for entropy,

$$\frac{H(X_1, X_2, \ldots, X_n)}{n} = \frac{\sum_{i=1}^{n} H(X_i|X^{i-1})}{n} \tag{4}$$
$$= \frac{H(X_n|X^{n-1}) + \sum_{i=1}^{n-1} H(X_i|X^{i-1})}{n} \tag{5}$$
$$= \frac{H(X_n|X^{n-1}) + H(X_1, X_2, \ldots, X_{n-1})}{n}. \tag{6}$$

From stationarity it follows that for all $1 \leq i \leq n$,

$$H(X_n|X^{n-1}) \leq H(X_i|X^{i-1}),$$

which further implies, by summing both sides over $i = 1, \ldots, n-1$ and dividing by $n-1$, that,

$$H(X_n|X^{n-1}) \leq \frac{\sum_{i=1}^{n-1} H(X_i|X^{i-1})}{n-1} \tag{7}$$
$$= \frac{H(X_1, X_2, \ldots, X_{n-1})}{n-1}. \tag{8}$$

Combining (6) and (8) yields,

$$\frac{H(X_1, X_2, \ldots, X_n)}{n} \leq \frac{1}{n}\left[\frac{H(X_1, X_2, \ldots, X_{n-1})}{n-1} + H(X_1, X_2, \ldots, X_{n-1})\right] \tag{9}$$
$$= \frac{H(X_1, X_2, \ldots, X_{n-1})}{n-1}. \tag{10}$$

Problem 6. By the chain rule for entropy,

$$H(X_0|X_{-1}, \ldots, X_{-n}) = H(X_0, X_{-1}, \ldots, X_{-n}) - H(X_{-1}, \ldots, X_{-n}) \tag{11}$$
$$= H(X_0, X_1, \ldots, X_n) - H(X_1, \ldots, X_n) \tag{12}$$
$$= H(X_0|X_1, \ldots, X_n), \tag{13}$$

where (12) follows from stationarity.

Problem 7. $X \multimap Y \multimap (Z, W)$ implies that $I(X; Z, W|Y) = 0$. Then,

$$I(X;Y) + I(Z;W) = I(X;Y) + I(X;Z,W|Y) + I(Z;W) = I(X;Y,Z,W) + I(Z;W)$$

Notice that $I(X;Y) + I(X;Z,W|Y) = I(X;Y,Z,W)$ follows from chain rule. Using the chain rule for a couple of times, we obtain the following steps.

$$I(X;Y,Z,W) + I(Z;W) = I(X;Z) + I(X;Y,W|Z) + I(Z;W) \tag{14}$$
$$= I(X;Z) + I(X;Y|W,Z) + I(X;W|Z) + I(Z;W) \tag{15}$$
$$= I(X;Z) + I(X;Y|W,Z) + I(X,Z;W) \tag{16}$$
$$\geq I(X;Z) + I(X;W) \tag{17}$$

$$\text{as } I(X,Z;W) \geq I(X;W)$$